

On intertwining operators and finite automorphism groups of vertex operator algebras

Kenichiro Tanabe*

Institute of Mathematics
University of Tsukuba

Abstract

Let V be a simple vertex operator algebra and G a finite automorphism group. We give a construction of intertwining operators for irreducible V^G -modules which occur as submodules of irreducible V -modules by using intertwining operators for V .

We also determine some fusion rules for a vertex operator algebra as an application.

1 Introduction

Let V be a simple vertex operator algebra (cf. [1], [8] and [9]), and G a finite automorphism group. It is an important problem to understand the module category for the vertex operator algebra V^G of G -invariants. In [2], this question was asked and several ideas are proposed.

For a simple vertex operator algebra V , it is shown in [6] that every irreducible V -module is a completely reducible V^G -module as a natural consequence of a duality theorem of Schur-Weyl type. In this paper we give a construction of intertwining operators for irreducible V^G -modules which occur as submodules of irreducible V -modules by using intertwining operators for V .

*Partially supported by JSPS Grant-in-Aid for Scientific Research No. 14740061.

Let's state our results more explicitly. Firstly, we need to recall the results of Dong–Yamshkulna[6]. For an irreducible V -module (L, Y_L) and $a \in G$ we define a new irreducible V -module $(L \circ a, Y_{L \circ a})$. Here $L \circ a$ is equal to L and $Y_{L \circ a}(u, z) = Y_L(au, z)$. Let $\mathcal{S}$ be a finite set of inequivalent irreducible V -modules which is closed under this right action of G . In [6] they define a finite dimensional semisimple associative algebra $\mathcal{A}_\alpha(G, \mathcal{S})$ over $\mathbb{C}$ and show a duality theorem of Schur-Weyl type for the actions of V^G and $\mathcal{A}_\alpha(G, \mathcal{S})$ on the direct sum of V -modules in $\mathcal{S}$ which is denoted by $\mathcal{L}$. That is, as a $\mathcal{A}_\alpha(G, \mathcal{S}) \otimes V^G$ -module, $\mathcal{L} = \bigoplus_{(j, \lambda) \in \Gamma} W_{(j, \lambda)} \otimes M_{(j, \lambda)}$, where $\{W_{(j, \lambda)}\}_{(j, \lambda) \in \Gamma}$ is the set of all inequivalent irreducible $\mathcal{A}_\alpha(G, \mathcal{S})$ -modules and $M_{(j, \lambda)}$ is the multiplicity spaces of $W_{(j, \lambda)}$ in $\mathcal{L}$. Each $M_{(j, \lambda)}$ is a nonzero irreducible V^G -module and the different multiplicity spaces are inequivalent V^G -modules. In this paper we consider intertwining operators for irreducible V^G -modules constructed from irreducible V -modules in this way.

For each $i = 1, 2, 3$ let $\mathcal{S}_i$ be a finite set of inequivalent irreducible V -modules which is closed under the action of G and let $\mathcal{L}_i$ be the direct sum of V -modules in $\mathcal{S}_i$. We have the decomposition $\mathcal{L}_i = \bigoplus_{(j_i, \lambda_i) \in \Gamma_i} W_{(j_i, \lambda_i)}^i \otimes M_{(j_i, \lambda_i)}^i$ as a $\mathcal{A}_{\alpha_i}(G, \mathcal{S}_i) \otimes V^G$ -module. Set $\mathcal{I} = \bigoplus_{(L^1, L^2, L^3) \in \mathcal{S}_1 \times \mathcal{S}_2 \times \mathcal{S}_3} I_V \left(\begin{smallmatrix} L^3 \\ L^1 & L^2 \end{smallmatrix} \right) \otimes L^1 \otimes L^2$, where $I_V \left(\begin{smallmatrix} L^3 \\ L^1 & L^2 \end{smallmatrix} \right)$ is the set of all intertwining operators of type $\left(\begin{smallmatrix} L^3 \\ L^1 & L^2 \end{smallmatrix} \right)$. $\mathcal{I}$ has a natural $\mathcal{A}_{\alpha_3}(G, \mathcal{S}_3)$ -module structure. For each $i = 1, 2, 3$, fix $(j_i, \lambda_i) \in \Gamma_i$ and nonzero $v^{10} \in M_{(j_1, \lambda_1)}^1, v^{20} \in M_{(j_2, \lambda_2)}^2$. Set $\mathcal{I}_{(j_1, \lambda_1), (j_2, \lambda_2)} = \text{Span}_{\mathbb{C}} \{ f \otimes (w^1 \otimes v^{10}) \otimes (w^2 \otimes v^{20}) \in \mathcal{I} \mid w^1 \in W_{(j_1, \lambda_1)}^1, w^2 \in W_{(j_2, \lambda_2)}^2 \}$. $\mathcal{I}_{(j_1, \lambda_1), (j_2, \lambda_2)}$ is a $\mathcal{A}_{\alpha_3}(G, \mathcal{S}_3)$ -submodule of $\mathcal{I}$. We will construct an injective linear map from the multiplicity space of $W_{(j_3, \lambda_3)}^3$ in $\mathcal{I}_{(j_1, \lambda_1), (j_2, \lambda_2)}$ to the set of all intertwining operators for V^G of type $\left(\begin{smallmatrix} M_{(j_3, \lambda_3)}^3 \\ M_{(j_1, \lambda_1)}^1 & M_{(j_2, \lambda_2)}^2 \end{smallmatrix} \right)$. Therefore, the fusion rule for V^G of type $\left(\begin{smallmatrix} M_{(j_3, \lambda_3)}^3 \\ M_{(j_1, \lambda_1)}^1 & M_{(j_2, \lambda_2)}^2 \end{smallmatrix} \right)$ is greater than or equal to the multiplicity of $W_{(j_3, \lambda_3)}^3$ in $\mathcal{I}_{(j_1, \lambda_1), (j_2, \lambda_2)}$.

This paper is organized as follows. In Sect.2 we first recall a construction of irreducible V^G -modules from irreducible V -modules in [6]. We also recall the definitions of intertwining operators and fusion rules. In Sect.3 we give a construction of intertwining operators for irreducible V^G -modules which occur as submodules of irreducible V -modules. In Sect.4 we apply the main result to determine some fusion rules for a vertex operator algebra $\mathcal{W}$ studied in [3]. In Appendix, we give some singular vectors in $\mathcal{W}$ -modules used in Sect. 4.

2 Preliminary

We assume that the reader is familiar with the basic knowledge on vertex operator algebras as presented in [1], [8], and [9].

The following notation will be in force throughout the paper: $V = (V, Y, \mathbf{1}, \omega)$ is a simple vertex operator algebra and G a finite automorphism group of V . For any V -module L , we always arrange the grading on $L = \bigoplus_{n=0}^{\infty} L(n)$ so that $L(0) \neq 0$ if $L \neq 0$ by using a grading shift.

2.1 Irreducible V^G -modules constructed from irreducible V -modules

In this subsection we review the results of Dong–Yamshkulna[6]. For a simple vertex operator algebra V , they showed that every irreducible V -module is a completely reducible V^G -module as a natural consequence of a duality theorem of Schur–Weyl type.

Let (L, Y_L) be an irreducible V -module and $a \in G$. We define a new irreducible V -module $(L \circ a, Y_{L \circ a})$. Here $L \circ a$ is equal to L and $Y_{L \circ a}(u, z) = Y_L(au, z)$. Note that $L \circ a$ is also an irreducible V -module. A set $\mathcal{S}$ of irreducible V -modules is called *stable* if for any $L \in \mathcal{S}$ and $a \in G$ there exists $M \in \mathcal{S}$ such that $L \circ a \simeq M$.

Now we take a finite G -stable set $\mathcal{S}$ consisting of inequivalent irreducible V -modules. Let $L \in \mathcal{S}$ and $a \in G$. Then there exists $M \in \mathcal{S}$ such that $M \simeq L \circ a^{-1}$. That is, there is a linear map $\phi(a, L) : L \rightarrow M$ satisfying the condition: $\phi(a, L)Y_L(v, z) = Y_M(av, z)\phi(a, L)$. By simplicity of L , there exists $\alpha_L(b, a) \in \mathbb{C}^\times$ such that $\phi(b, M)\phi(a, L) = \alpha_L(b, a)\phi(ba, L)$. Moreover, for $a, b, c \in G$ we have

$$\alpha_L(c, ba)\alpha_L(b, a) = \alpha_M(c, b)\alpha_L(cb, a).$$

For $L \in \mathcal{S}$ and $a \in G$, we denote $M \in \mathcal{S}$ such that $L \circ a \simeq M$ by $L \cdot a$.

Define a vector space $\mathbb{CS} = \bigoplus_{L \in \mathcal{S}} \mathbb{C}e(L)$ with a basis $e(L)$ for $L \in \mathcal{S}$. The space $\mathbb{CS}$ is an associative algebra under the product $e(L)e(M) = \delta_{L,M}e(L)$. Let $\mathcal{U}(\mathbb{CS}) = \{\sum_{L \in \mathcal{S}} \lambda_L e(L) \mid \lambda_L \in \mathbb{C}^\times\}$ be the set of unit elements on $\mathbb{CS}$. $\mathcal{U}(\mathbb{CS})$ is a multiplicative right G -set by the action $(\sum_{L \in \mathcal{S}} \lambda_L e(L)) \cdot a = \sum_{L \in \mathcal{S}} \lambda_L e(L \cdot a)$ for $a \in G$. Set $\alpha(a, b) = \sum_{L \in \mathcal{S}} \alpha_L(a, b)e(L)$. Then $(\alpha(a, b) \cdot c)\alpha(ab, c) = \alpha(a, bc)\alpha(b, c)$ hold for all $a, b, c \in G$. So $\alpha : G \times G \rightarrow \mathcal{U}(\mathbb{CS})$ is a 2-cocycle.

Define the vector space $\mathcal{A}_\alpha(G, \mathcal{S}) = \mathbb{C}[G] \otimes \mathbb{C}\mathcal{S}$ with a basis $a \otimes e(L)$ for $a \in G$ and $L \in \mathcal{S}$ and a multiplication on it:

$$a \otimes e(L) \cdot b \otimes e(M) = \alpha_M(a, b)ab \otimes e(L \cdot b)e(M).$$

Then $\mathcal{A}_\alpha(G, \mathcal{S})$ is an associative algebra with the identity element $\sum_{L \in \mathcal{S}} 1 \otimes e(L)$.

We define an action of $\mathcal{A}_\alpha(G, \mathcal{S})$ on $\mathcal{L} = \bigoplus_{L \in \mathcal{S}} L$ as follows: For $L, M \in \mathcal{S}$, $w \in M$ and $a \in G$ we set

$$a \otimes e(L) \cdot w = \delta_{L,M} \phi(a, L)w.$$

Note that the actions of $\mathcal{A}_\alpha(G, \mathcal{S})$ and V^G on $\mathcal{L}$ commute with each other.

For each $L \in \mathcal{S}$ set $G_L = \{a \in G \mid L \circ a \simeq L \text{ as } V\text{-modules}\}$. Let $\mathcal{O}_L$ be the orbit of L under the action of G and let $G = \sqcup_{j=1}^k G_L g_j$ be a right coset decomposition with $g_1 = 1$. Then $\mathcal{O}_L = \{L \cdot g_j \mid j = 1, \dots, k\}$ and $G_{L \cdot g_j} = g_j^{-1} G_L g_j$. We define several subspaces of $\mathcal{A}_\alpha(G, \mathcal{S})$ by:

$$\begin{aligned} S(L) &= \text{Span}_{\mathbb{C}}\{a \otimes e(L) \mid a \in G_L\}, \\ D(L) &= \text{Span}_{\mathbb{C}}\{a \otimes e(L) \mid a \in G\} \text{ and} \\ D(\mathcal{O}_L) &= \text{Span}_{\mathbb{C}}\{a \otimes e(L \cdot g_j) \mid j = 1, \dots, k, a \in G\}. \end{aligned}$$

Decompose $\mathcal{S}$ into a disjoint union of orbits $\mathcal{S} = \sqcup_{j \in J} \mathcal{O}_j$. Let $L^{(j)}$ be a representative element of $\mathcal{O}_j$. Then $\mathcal{O}_j = \{L^{(j)} \cdot a \mid a \in G\}$ and $\mathcal{A}_\alpha(G, \mathcal{S}) = \bigoplus_{j \in J} D(\mathcal{O}_{L^{(j)}})$. We recall the following properties of $\mathcal{A}_\alpha(G, \mathcal{S})$.

Lemma 1. ([6], Lemma 3.4) *Let $L \in \mathcal{S}$ and $G = \sqcup_{j=1}^k G_L g_j$. Then*

- (1) *$S(L)$ is a subalgebra of $\mathcal{A}_\alpha(G, \mathcal{S})$ isomorphic to $\mathbb{C}^{\alpha_L}[G_L]$ where $\mathbb{C}^{\alpha_L}[G_L]$ is the twisted group algebra with 2-cocycle α_L .*
- (2) *$D(\mathcal{O}_L) = \bigoplus_{j=1}^k D(L \cdot g_j)$ is a direct sum of left ideals.*
- (3) *Each $D(\mathcal{O}_L)$ is a two sided ideal of $\mathcal{A}_\alpha(G, \mathcal{S})$ and $\mathcal{A}_\alpha(G, \mathcal{S}) = \bigoplus_{j \in J} D(\mathcal{O}_{L^{(j)}})$. Moreover, $D(\mathcal{O}_L)$ has the identity element $\sum_{M \in \mathcal{O}_L} 1 \otimes e(M)$.*

Lemma 2. ([6], Theorem 3.6)

- (1) *$D(\mathcal{O}_L)$ is semisimple for all $L \in \mathcal{S}$ and the simple $D(\mathcal{O}_L)$ -modules are precisely equal to $\text{Ind}_{S(L)}^{D(L)} U = D(L) \otimes_{S(L)} U$ where U ranges over the simple $\mathbb{C}^{\alpha_L}[G_L]$ -modules.*

(2) $\mathcal{A}_\alpha(G, \mathcal{S})$ is semisimple and simple $\mathcal{A}_\alpha(G, \mathcal{S})$ -modules are precisely $\text{Ind}_{S(L^{(j)})}^{D(L^{(j)})} U$ where U ranges over the simple $\mathbb{C}^{\alpha_{L^{(j)}}}[G_{L^{(j)}}]$ -modules and $j \in J$.

For $L \in \mathcal{S}$ let Λ_{G_L, α_L} be the set of all irreducible characters λ of $\mathbb{C}^{\alpha_L}[G_L]$. We denote the corresponding simple module by $U(L, \lambda)$. Note that L is a semisimple $\mathbb{C}^{\alpha_L}[G_L]$ -module. Let L^λ be the sum of simple $\mathbb{C}^{\alpha_L}[G_L]$ -submodules of L isomorphic to $U(L, \lambda)$. Then $L = \bigoplus_{\lambda \in \Lambda_{G_L, \alpha_L}} L^\lambda$. Moreover $L^\lambda = U(L, \lambda) \otimes L_\lambda$ where $L_\lambda = \text{Hom}_{\mathbb{C}^{\alpha_L}[G_L]}(U(L, \lambda), L)$ is the multiplicity space of $U(L, \lambda)$ in L . We can realize L_λ as a subspace of L in the following way: Let $w \in U(L, \lambda)$ be a fixed nonzero vector. Then we can identify $\text{Hom}_{\mathbb{C}^{\alpha_L}[G_L]}(U(L, \lambda), L)$ with the subspace

$$\{f(w) \mid f \in \text{Hom}_{\mathbb{C}^{\alpha_L}[G_L]}(U(L, \lambda), L)\}$$

of L^λ . Note that the actions of $\mathbb{C}^{\alpha_L}[G_L]$ and V^{G_L} on L commute with each other. So L^λ and L_λ are ordinary V^{G_L} -modules. Furthermore, L^λ and L_λ are ordinary V^G -modules.

For convenience, we set $G_j = G_{L^{(j)}}$, $\Lambda_j = \Lambda_{L^{(j)}, \alpha_{L^{(j)}}}$ and $U_{(j, \lambda)} = U(L^{(j)}, \lambda)$ for $j \in J$ and $\lambda \in \Lambda_j$. We denote by Γ the set $\{(j, \lambda) \mid j \in J, \lambda \in \Lambda_j\}$. We have a decomposition

$$L^{(j)} = \bigoplus_{\lambda \in \Lambda_j} U_{(j, \lambda)} \otimes M_{(j, \lambda)}$$

as a $\mathbb{C}^{\alpha_{L^{(j)}}}[G_j] \otimes V^{G_j}$ -module. We also have

$$\mathcal{L} = \bigoplus_{(j, \lambda) \in \Gamma} \text{Ind}_{S(L^{(j)})}^{D(L^{(j)})} U_{(j, \lambda)} \otimes M_{(j, \lambda)}$$

as a $\mathcal{A}_\alpha(G, \mathcal{S}) \otimes V^G$ -module.

For $(j, \lambda) \in \Gamma$ we set $W_{(j, \lambda)} = \text{Ind}_{S(L^{(j)})}^{D(L^{(j)})} U_{(j, \lambda)}$. Then $W_{(j, \lambda)}$ forms a complete list of simple $\mathcal{A}_\alpha(G, \mathcal{S})$ -modules by Lemma 2.

A duality theorem of Schur–Weyl type holds.

Theorem 1. ([6] Theorem 6.14) *As a $\mathcal{A}_\alpha(G, \mathcal{S}) \otimes V^G$ -module,*

$$\mathcal{L} = \bigoplus_{(j, \lambda) \in \Gamma} W_{(j, \lambda)} \otimes M_{(j, \lambda)}.$$

Moreover,

- (1) Each $M_{(j,\lambda)}$ is a nonzero irreducible V^G -module.
- (2) $M_{(j_1,\lambda_1)}$ and $M_{(j_2,\lambda_2)}$ are isomorphic V^G -modules if and only if $(j_1, \lambda_1) = (j_2, \lambda_2)$.

2.2 Intertwining operators and fusion rules

We recall the definition of intertwining operators for V -modules which is introduced in [8].

Definition 1. Let L^i ($i = 1, 2, 3$) be V -modules. An intertwining operator of type $\begin{pmatrix} L^3 \\ L^1 \ L^2 \end{pmatrix}$ is a linear map

$$\begin{aligned} I(\cdot, z) : L^1 &\rightarrow \text{Hom}_{\mathbb{C}}(L^2, L^3)\{z\}, \\ v &\mapsto I(v, z) = \sum_{\gamma \in \mathbb{C}} v_{\gamma}^{-\gamma-1}, \end{aligned}$$

which satisfies the following conditions: Let $u \in V, v \in L^1$, and $w \in L^2$.

(1) For any fixed $\gamma \in \mathbb{C}$, $v_{\gamma+n}w = 0$ for $n \in \mathbb{Z}$ sufficiently large.

$$(2) \quad I(L(-1)v, z) = \frac{d}{dz}I(v, z).$$

(3)

$$\begin{aligned} & z_0^{-1} \delta\left(\frac{z_1 - z_2}{z_0}\right) Y_{L^3}(u, z_1) I(v, z_2) w - z_0^{-1} \delta\left(\frac{-z_2 + z_1}{z_0}\right) I(v, z_2) Y_{L^2}(u, z_1) w \\ &= z_2^{-1} \delta\left(\frac{z_1 - z_0}{z_2}\right) I(Y_{L^1}(u, z_0) v, z_2) w. \end{aligned}$$

We denote by $I_V\left(\begin{smallmatrix} L^3 \\ L^1 \ L^2 \end{smallmatrix}\right)$ the set of all intertwining operators of type $\begin{pmatrix} L^3 \\ L^1 \ L^2 \end{pmatrix}$. The dimension of $I_V\left(\begin{smallmatrix} L^3 \\ L^1 \ L^2 \end{smallmatrix}\right)$ is called the *fusion rule* of type $\begin{pmatrix} L^3 \\ L^1 \ L^2 \end{pmatrix}$.

For a V -module $L = \bigoplus_{n=0}^{\infty} L(n)$, it is shown in [8, Theorem 5.2.1.] that the graded vector space $L' = \bigoplus_{n=0}^{\infty} L(n)^*$ carries the structure of a V -module, where $L(n)^* = \text{Hom}_{\mathbb{C}}(L(n), \mathbb{C})$. L' is called the *contragredient module* of L .

The fusion rules have some symmetries.

Lemma 3. ([8], Proposition 5.4.7 and 5.5.2) *Let L^i ($i = 1, 2, 3$) be V -modules. Then*

$$\dim_{\mathbb{C}} I_V\left(\begin{smallmatrix} L^3 \\ L^1 \ L^2 \end{smallmatrix}\right) = \dim_{\mathbb{C}} I_V\left(\begin{smallmatrix} L^3 \\ L^2 \ L^1 \end{smallmatrix}\right) = \dim_{\mathbb{C}} I_V\left(\begin{smallmatrix} (L^2)' \\ L^1 \ (L^3)' \end{smallmatrix}\right).$$

Let L^1 and L^2 be irreducible V -modules. We use a notation $L^1 \times L^2 = \sum_{L^3} \dim_{\mathbb{C}} I_V \left(\begin{smallmatrix} L^3 \\ L^1 \ L^2 \end{smallmatrix} \right) L^3$ to represent the fusion rules, where L^3 ranges over the irreducible V -modules. Note that $L^1 \times L^2 = L^2 \times L^1$ by Lemma 3.

3 Intertwining operators for irreducible V^G -modules which occur as submodules of irreducible V -modules

In this section we give a construction of intertwining operators for irreducible V^G -modules which occur as submodules of irreducible V -modules by using intertwining operators for V .

Let $\mathcal{S}_i$ ($i = 1, 2, 3$) be finite G -stable sets consisting of inequivalent irreducible V -modules. Set $\mathcal{L}_i = \bigoplus_{L \in \mathcal{S}_i} L$. For $L^i \in \mathcal{S}_i$ and $a \in G$, $\phi_i(a, L^i) : L^i \rightarrow L^i \cdot a^{-1}$ denote the fixed V -module isomorphisms. For $L^i \in \mathcal{S}_i$ and $a, b \in G$, $\alpha_{i, L^i} \in \mathbb{C}^\times$ denote nonzero complex numbers such that $\phi_i(b, L^i \cdot a^{-1}) \phi_i(a, L^i) = \alpha_{i, L^i}(b, a) \phi(ba, L^i)$. Set $\alpha_i(a, b) = \sum_{L^i \in \mathcal{S}_i} \alpha_{i, L^i}(a, b) e(L^i)$. Let $\mathcal{S}_i = \sqcup_{j \in J_i} \mathcal{O}_j$ be the orbit decompositions under the action of G and set $\Gamma_i = \{(j_i, \lambda_i) \mid j_i \in J_i, \lambda_i \in \Lambda_{j_i}\}$.

For $f \in I_V \left(\begin{smallmatrix} L^3 \\ L^1 \ L^2 \end{smallmatrix} \right)$ and $a \in G$, we define ${}_a f \in I_V \left(\begin{smallmatrix} L^3 \cdot a^{-1} \\ L^1 \cdot a^{-1} \ L^2 \cdot a^{-1} \end{smallmatrix} \right)$ as follows: For $v \in L^1$ we set

$${}_a f(v, z) = \phi_3(a, L^3) f(\phi_1(a, L^1)^{-1} v, z) \phi_2(a, L^2)^{-1}.$$

Set

$$\mathcal{I} = \bigoplus_{(L^1, L^2, L^3) \in \mathcal{S}_1 \times \mathcal{S}_2 \times \mathcal{S}_3} I_V \left(\begin{smallmatrix} L^3 \\ L^1 \ L^2 \end{smallmatrix} \right) \otimes_{\mathbb{C}} L^1 \otimes_{\mathbb{C}} L^2.$$

We define an action of $\mathcal{A}_{\alpha_3}(G, \mathcal{S})$ on $\mathcal{I}$ as follows: Let $L^i \in \mathcal{S}_i$ ($i = 1, 2, 3$). For $a \otimes e(M) \in \mathcal{A}_{\alpha_3}(G, \mathcal{S})$, $v \in L^1$, $w \in L^2$, and $f \in I_V \left(\begin{smallmatrix} L^3 \\ L^1 \ L^2 \end{smallmatrix} \right)$, we set

$$\begin{aligned} (a \otimes e(M)) \cdot (f \otimes v \otimes w) &= \delta_{M, L^3} \cdot {}_a f \otimes \phi_1(a, L^1) v \otimes \phi_2(a, L^2) w \\ &\in I_V \left(\begin{smallmatrix} L^3 \cdot a^{-1} \\ L^1 \cdot a^{-1} \ L^2 \cdot a^{-1} \end{smallmatrix} \right) \otimes L^1 \cdot a^{-1} \otimes L^2 \cdot a^{-1}. \end{aligned}$$

We define a map $\Psi : \mathcal{I} \rightarrow \mathcal{L}_3\{z\}$ by

$$\Psi(f \otimes v^1 \otimes v^2) = f(v^1, z) v^2$$

for $v^1 \in L^1$, $v^2 \in L^2$, and $f \in I_V\left(\begin{smallmatrix} L^3 \\ L^1 \ L^2 \end{smallmatrix}\right)$, where $L^i \in \mathcal{S}_i$ ($i = 1, 2, 3$). Note that Ψ is a $\mathcal{A}_{\alpha_3}(G, \mathcal{S}_3)$ -module homomorphism.

Lemma 4. *The map $\Psi : \mathcal{I} \rightarrow \mathcal{L}_3\{z\}$ is injective.*

Proof. We use the same method that was used in the proof of Lemma 3.1. of [5]. Assume false. Then there is a nonzero $X \in \text{Ker}\Psi$. Since $\mathcal{L}_3 = \oplus_{L \in \mathcal{S}_3} L$, we may assume $X = \sum_{i,j} f^{ij} \otimes v^{1i} \otimes v^{2j}$, where $v^{1i} \in L^{1i}$ ($i = 1, \dots, l_1$) are linearly independent homogeneous vectors in $\mathcal{L}_1$, $v^{2j} \in L^{2j}$ ($j = 1, \dots, l_2$) are linearly independent homogeneous vectors in $\mathcal{L}_2$, $f^{ij} \in I_V\left(\begin{smallmatrix} L^3 \\ L^{1i} \ L^{2j} \end{smallmatrix}\right)$, $L^{1i} \in \mathcal{S}_1$, $L^{2j} \in \mathcal{S}_2$, and $L^3 \in \mathcal{S}_3$. We may also assume $f^{11} \otimes v^{11} \otimes v^{21}$ is nonzero. Since $\sum_{i,j} f^{ij}(v^{1i}, z)v^{2j} = 0$, for all $u \in V$ we have

$$\sum_{i,j} Y_{\mathcal{L}_3}(u, z_1) f^{ij}(v^{1i}, z) v^{2j} = 0.$$

Using the associativity of intertwining operators [4, Proposition 11.5],

$$\sum_{i,j} f^{ij}(Y_{\mathcal{L}_1}(u, z_1) v^{1i}, z) v^{2j} = 0. \quad (1)$$

We denote $Y_{\mathcal{L}_1}(u, z_1) = \sum_{n \in \mathbb{Z}} u_n^{\mathcal{L}_1} z_1^{-n-1}$. Fix $N \in \mathbb{Z}$ such that $v^{1i} \in \oplus_{n=0}^N L^{1i}(n)$ for all i . Since $\mathcal{S}_1$ consists of inequivalent irreducible V -modules, the linear map $\sigma_N : V \rightarrow \oplus_{L \in \mathcal{S}_1} \oplus_{m=0}^N \text{End}_{\mathbb{C}} L(m)$ defined by $\sigma_N(u) = u_{\text{wt}u-1}^{\mathcal{L}_1}$ for homogeneous $u \in V$ is an epimorphism by Lemma 6.13 of [6]. So there exists $u^1 \in V$ such that $\sigma_N(u^1) v^{1i} = \delta_{1,i} v^{11}$. From formula (1), we have

$$0 = \sum_{i,j} f^{ij}(\sigma_N(u^1) v^{1i}, z) v^{2j} = \sum_j f^{1j}(v^{11}, z) v^{2j}. \quad (2)$$

Therefore, for all $u \in V$ we have

$$\sum_j Y_{\mathcal{L}_3}(u, z_1) f^{1j}(v^{11}, z) v^{2j} = 0.$$

Using the commutativity of intertwining operators [4, Proposition 11.4],

$$\sum_j f^{1j}(v^{11}, z) Y_{\mathcal{L}_2}(u, z_1) v^{2j} = 0.$$

Since $\mathcal{S}_2$ consists of inequivalent irreducible V -modules, we have

$$f^{11}(v^{11}, z)v^{21} = 0$$

for the same reason to obtain formula (2). So $f^{11} \otimes v^{11} \otimes v^{21} \in \text{Ker } \Psi$. Since L^{11} and L^{21} are irreducible V -modules and $f^{11} \otimes v^{11} \otimes v^{21}$ is nonzero, this contradicts Proposition 11.9 of [4]. $\square$

We have the decomposition of each $\mathcal{L}_i$ as a $\mathcal{A}_{\alpha_i}(G, \mathcal{S}_i)$ -module in Theorem 1:

$$\mathcal{L}_i = \bigoplus_{(j_i, \lambda_i) \in \Gamma_i} W_{(j_i, \lambda_i)}^i \otimes M_{(j_i, \lambda_i)}^i.$$

For $i = 1, 2$ let $(j_i, \lambda_i) \in \Gamma_i$ and let $v^{i0} \in M_{(j_i, \lambda_i)}^i$. Set

$$\begin{aligned} & \mathcal{I}_{(j_1, \lambda_1), (j_2, \lambda_2)}(v^{10}, v^{20}) \\ &= \text{Span}_{\mathbb{C}} \{f \otimes (w^1 \otimes v^{10}) \otimes (w^2 \otimes v^{20}) \in \mathcal{I} \mid w^1 \in W_{(j_1, \lambda_1)}^1, w^2 \in W_{(j_2, \lambda_2)}^2\}. \end{aligned}$$

$\mathcal{I}_{(j_1, \lambda_1), (j_2, \lambda_2)}(v^{10}, v^{20})$ is a $\mathcal{A}_{\alpha_3}(G, \mathcal{S}_3)$ -submodule of $\mathcal{I}$. For any nonzero $v^{10}, v^1 \in M_{(j_1, \lambda_1)}^1$ and nonzero $v^{20}, v^2 \in M_{(j_2, \lambda_2)}^2$, $\mathcal{I}_{(j_1, \lambda_1), (j_2, \lambda_2)}(v^{10}, v^{20})$ and $\mathcal{I}_{(j_1, \lambda_1), (j_2, \lambda_2)}(v^1, v^2)$ are isomorphic $\mathcal{A}_{\alpha_3}(G, \mathcal{S}_3)$ -modules.

Theorem 2. Fix a nonzero $v^{10} \in M_{(j_1, \lambda_1)}^1$ and a nonzero $v^{20} \in M_{(j_2, \lambda_2)}^2$.

For any $((j_1, \lambda_1), (j_2, \lambda_2), (j_3, \lambda_3)) \in \Gamma_1 \times \Gamma_2 \times \Gamma_3$, there exists an injective linear map from $\text{Hom}_{\mathcal{A}_{\alpha_3}(G, \mathcal{S}_3)}(W_{(j_3, \lambda_3)}^3, \mathcal{I}_{(j_1, \lambda_1), (j_2, \lambda_2)}(v^{10}, v^{20}))$ to $I_{VG} \left(\begin{smallmatrix} M_{(j_3, \lambda_3)}^3 \\ M_{(j_1, \lambda_1)}^1 \ M_{(j_2, \lambda_2)}^2 \end{smallmatrix} \right)$. In particular,

$$\dim_{\mathbb{C}} I_{VG} \left(\begin{smallmatrix} M_{(j_3, \lambda_3)}^3 \\ M_{(j_1, \lambda_1)}^1 \ M_{(j_2, \lambda_2)}^2 \end{smallmatrix} \right) \geq \dim_{\mathbb{C}} \text{Hom}_{\mathcal{A}_{\alpha_3}(G, \mathcal{S}_3)}(W_{(j_3, \lambda_3)}^3, \mathcal{I}_{(j_1, \lambda_1), (j_2, \lambda_2)}(v^{10}, v^{20})).$$

Proof. For convenience, we set $W^i = W_{(j_i, \lambda_i)}^i$, $M^i = M_{(j_i, \lambda_i)}^i$ ($i = 1, 2, 3$), $\mathcal{A}_3 = \mathcal{A}_{\alpha_3}(G, \mathcal{S}_3)$, and $\mathcal{I}_{12}(v^{10}, v^{20}) = \mathcal{I}_{(j_1, \lambda_1), (j_2, \lambda_2)}(v^{10}, v^{20})$. Fix a nonzero $w^{30} \in W^3$. Let $F \in \text{Hom}_{\mathcal{A}_3}(W^3, \mathcal{I}_{12}(v^{10}, v^{20}))$. We shall define $\Phi(F) \in I_{VG} \left(\begin{smallmatrix} M^3 \\ M^1 \ M^2 \end{smallmatrix} \right)$.

For $v^1 \in M^1, v^2 \in M^2$, we set

$$H(v^1, v^2) \in \text{Hom}_{\mathcal{A}_3}(W^3, \mathcal{I}_{12}(v^1, v^2))$$

as follows: For $w^3 \in W^3$, let $F(w^3) = \sum_i f^i \otimes (w^{1i} \otimes v^{10}) \otimes (w^{2i} \otimes v^{20}) \in \mathcal{I}_{12}(v^{10}, v^{20})$, where $w^{1i} \otimes v^{10} \in L^{1i}, w^{2i} \otimes v^{20} \in L^{2i}, f^i \in I_V \left(\begin{smallmatrix} L^{3i} \\ L^{1i} \ L^{2i} \end{smallmatrix} \right)$ and

$L^{ji} \in \mathcal{S}_j$ ($j = 1, 2, 3$). Note that $w^{ji} \otimes v^j \in L^{ji}$ by the definition of M^j for $j = 1, 2$. We define

$$H(v^1, v^2)(w^3) = \sum_i f^i \otimes (w^{1i} \otimes v^1) \otimes (w^{2i} \otimes v^2)$$

It is clear that $H(v^1, v^2) \in \text{Hom}_{\mathcal{A}_3}(W^3, \mathcal{I}_{12}(v^1, v^2))$. Since the map $\Psi : \mathcal{I} \rightarrow \mathcal{L}_3\{z\}$ is a $\mathcal{A}_3$ -module homomorphism, the map $w^3 \mapsto \Psi(H(v^1, v^2)(w^3))$ is a $\mathcal{A}_3$ -module homomorphism from W^3 to $\mathcal{L}_3\{z\}$. So $\Psi(H(v^1, v^2)(W^3))$ is a $\mathcal{A}_3$ -submodule of $(W^3 \otimes M^3)\{z\}$. Let $w^{3,1}, w^{3,2}, \dots, w^{3, \dim_{\mathbb{C}} W^3}$ be a basis of W^3 . Since W^3 is an irreducible $\mathcal{A}_3$ -module, there exists $a \in \mathcal{A}_3$ such that $aw^{3,i} = \delta_{1,i}w^{3,1}$. Let $\Psi(H(v^1, v^2)(w^{3,1})) = \sum_i w^{3,i} \otimes p^i$, where $p^i \in M^3\{z\}$. Then

$$\begin{aligned} \Psi(H(v^1, v^2)(w^{3,1})) &= \Psi(H(v^1, v^2)(aw^{3,1})) = a\Psi(H(v^1, v^2)(w^{3,1})) \\ &= a \sum_i w^{3,i} \otimes p^i = \sum_i (aw^{3,i}) \otimes p^i = w^{3,1} \otimes p^1. \end{aligned}$$

So $\Psi(H(v^1, v^2)(w^3)) \in (w^3 \otimes M^3)\{z\}$ for all $w^3 \in W^3$. We hence have an unique $\Phi(F)(v^1, z)v^2 \in M^3\{z\}$ such that

$$w^{30} \otimes \Phi(F)(v^1, z)v^2 = \Psi(H(v^1, v^2)(w^{30})). \quad (3)$$

Since f^i are intertwining operators, we have $\Phi(F) \in I_{VG} \left(\begin{smallmatrix} M^3 \\ M^1 & M^2 \end{smallmatrix} \right)$ from formula (3).

We will show that Φ is injective. Suppose $\Phi(F) = 0$. Then

$$0 = w^{30} \otimes \Phi(F)(v^{10}, z)v^{20} = \Psi(F(w^{30})).$$

Since Ψ is injective by Lemma 4, $F(w^{30}) = 0$. Since W^3 is an irreducible $\mathcal{A}_3$ -module and $F \in \text{Hom}_{\mathcal{A}_3}(W^3, \mathcal{I}_{12}(v^{10}, v^{20}))$, $F = 0$. $\square$

Let $\mathbb{C}G$ be the group algebra of G and $\text{Irr}G$ the set of all irreducible characters of G . We set $\mathcal{S}_i = \{V\}$ ($i = 1, 2, 3$) in Theorem 2. Then $\mathcal{A}_{\alpha_i}(G, \mathcal{S}_i) = \mathbb{C}G$ and $\Gamma_i = \text{Irr}G$. Note that $\dim_{\mathbb{C}} I_V \left(\begin{smallmatrix} V \\ V & V \end{smallmatrix} \right) = 1$ since V is simple. In this case we have the following result:

Corollary 3. *Let $\chi_i \in \text{Irr}G$ ($i = 1, 2, 3$). Then*

$$\dim_{\mathbb{C}} I_{VG} \left(\begin{smallmatrix} V_{\chi_3} \\ V_{\chi_1} & V_{\chi_2} \end{smallmatrix} \right) \geq \dim_{\mathbb{C}} \text{Hom}_{\mathbb{C}G}(W_{\chi_3}, W_{\chi_1} \otimes_{\mathbb{C}} W_{\chi_2}).$$

In [2, Section 3], it is conjectured that if V is rational then for all $\chi_1, \chi_2 \in \text{Irr}G$,

$$V_{\chi_1} \times V_{\chi_2} = \sum_{\chi_3 \in \text{Irr}G} \dim_{\mathbb{C}} \text{Hom}_{\mathbb{C}G}(W_{\chi_3}, W_{\chi_1} \otimes_{\mathbb{C}} W_{\chi_2}) V_{\chi_3}.$$

The conjecture implies that if V is rational then the representation algebra of the finite group G is always realized as a subalgebra of the fusion algebra of V^G .

4 An application

In [3] we studied a vertex operator algebra $\mathcal{W}$ which is a realization of an algebra denoted by $[Z_3^{(5)}]$ in [7]. $\mathcal{W}$ is a fixed point subalgebra of a vertex operator algebra M_k^0 . It is expected that the $\mathbb{Z}_3$ symmetry of $\mathcal{W}$ affords 3B elements of the Monster simple group [11]. For M_k^0 , the irreducible modules are classified and the fusion rules are determined in [12]. Let L^i ($i = 1, 2, 3$) be irreducible $\mathcal{W}$ -modules such that L^1 and L^2 occur as submodules of irreducible M_k^0 -modules. In this section we determine the fusion rule of type $\begin{pmatrix} L^3 \\ L^1 \ L^2 \end{pmatrix}$ by using Theorem 2.

4.1 Subalgebra M_k^0 of $V_{\sqrt{2}A_2}$

In this subsection we review some properties of M_k^0 in [12]. Let A_2 be the ordinary root lattice of type A_2 and $V_{\sqrt{2}A_2}$ the lattice vertex operator algebra associated with $\sqrt{2}A_2$. Let α_1, α_2 be the simple roots of type A_2 and set $\alpha_0 = -\alpha_1 - \alpha_2$.

For basic definitions concerning lattice vertex operator algebras we refer to [4] and [9]. Our notation for the lattice vertex operator algebra is standard [9]. In particular, $\mathfrak{h} = \mathbb{C} \otimes_{\mathbb{Z}} \sqrt{2}A_2$ is an abelian Lie algebra, $\hat{\mathfrak{h}} = \mathfrak{h} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c$ is the corresponding affine Lie algebra, $M(1) = \mathbb{C}[\alpha(n) ; \alpha \in \mathfrak{h}, n < 0]$, where $\alpha(n) = \alpha \otimes t^n$, is the unique irreducible $\hat{\mathfrak{h}}$ -module such that $\alpha(n)1 = 0$ for all $\alpha \in \mathfrak{h}, n > 0$, and $c = 1$. As a vector space $V_{\sqrt{2}A_2} = M(1) \otimes \mathbb{C}[\sqrt{2}A_2]$ and for each $v \in V_{\sqrt{2}A_2}$, a vertex operator $Y(v, z) = \sum_{n \in \mathbb{Z}} v_n z^{-n-1} \in \text{End}(V_{\sqrt{2}A_2})[[z, z^{-1}]]$ is defined. The vector $1 = 1 \otimes 1$ is called the vacuum vector. We use the symbol $e^\alpha, \alpha \in \sqrt{2}A_2$ to denote a basis of $\mathbb{C}[\sqrt{2}A_2]$.

There exists an isometry τ of $\sqrt{2}A_2$ such that $\tau(\sqrt{2}\alpha_1) = \sqrt{2}\alpha_2$ and $\tau(\sqrt{2}\alpha_2) = \sqrt{2}\alpha_0$. The isometry τ lifts naturally to an automorphism of $V_{\sqrt{2}A_2}$:

$$\alpha^1(-n_1) \cdots \alpha^k(-n_k)e^\beta \mapsto (\tau\alpha^1)(-n_1) \cdots (\tau\alpha^k)(-n_k)e^{\tau\beta}.$$

By abuse of notation, we denote it by τ . Let G be the cyclic group generated by τ . Set

$$\begin{aligned} \omega^3 &= \frac{1}{15}(\alpha_1(-1)^2 + \alpha_2(-1)^2 + \alpha_0(-1)^2) \\ &\quad + \frac{1}{10}(e^{\sqrt{2}\alpha_1} + e^{-\sqrt{2}\alpha_1} + e^{\sqrt{2}\alpha_2} + e^{-\sqrt{2}\alpha_2} + e^{\sqrt{2}\alpha_0} + e^{-\sqrt{2}\alpha_0}) \end{aligned}$$

and $M_k^0 = \{v \in V_{\sqrt{2}A_2} \mid (\omega^3)_1 v = 0\}$. Since $\tau\omega^3 = \omega^3$, M_k^0 is invariant under the action of τ . τ is an automorphism group of M_k^0 of order 3 by [3, Theorem 2.1]. Set

$$L_0 = L, \quad L_a = \frac{\sqrt{2}\alpha_2}{2} + L, \quad L_b = \frac{\sqrt{2}\alpha_0}{2} + L, \quad L_c = \frac{\sqrt{2}\alpha_1}{2} + L$$

and

$$\begin{aligned} M_k^i &= \{v \in V_{L_i} \mid (\omega^3)_1 v = 0\}, \\ W_k^i &= \{v \in V_{L_i} \mid (\omega^3)_1 v = \frac{2}{5}v\}, \quad \text{for } i = 0, a, b, c. \end{aligned}$$

It is shown in [12] that $\{M_k^i, W_k^i \mid i = 0, a, b, c\}$ is the set of all irreducible M_k^0 -modules and the fusion rules are determined.

4.2 Subalgebra $\mathcal{W}$ in M^τ

We denote by $\mathcal{W}$ the subalgebra $(M_k^0)^\tau$ of fixed points of τ in M_k^0 . We recall some properties of $\mathcal{W}$ in [3]. $\mathcal{W}$ is generated by the Virasoro element ω and an element J of weight 3. Let $Y(\omega, z) = \sum_{n \in \mathbb{Z}} L(n)z^{-n-2}$ and $Y(J, z) =$

$\sum_{n \in \mathbb{Z}} J(n)z^{-n-3}$. They satisfy the following commutation relations:

$$\begin{aligned}
[L(m), L(n)] &= (m-n)L(m+n) + \frac{m^3-m}{12} \cdot \frac{6}{5} \cdot \delta_{m+n,0}, \\
[L(m), J(n)] &= (2m-n)J(m+n), \\
[J(m), J(n)] &= (m-n)(22(m+n+2)(m+n+3) + 35(m+2)(n+2))L(m+n) \\
&\quad - 120(m-n) \left(\sum_{k \leq -2} L(k)L(m+n-k) + \sum_{k \geq -1} L(m+n-k)L(k) \right) \\
&\quad - \frac{7}{10}m(m^2-1)(m^2-4)\delta_{m+n,0}.
\end{aligned} \tag{4}$$

$\mathcal{W}$ has exactly 20 irreducible modules. 8 irreducible $\mathcal{W}$ -modules occur as submodules of irreducible M_k^0 -modules. We introduce those 8 $\mathcal{W}$ -modules. τ acts on the irreducible M_k^0 -modules as follows:

$$\begin{aligned}
M_k^0 \circ \tau &\simeq M_k^0, W_k^0 \circ \tau \simeq W_k^0, \\
M_k^a \circ \tau &\simeq M_k^c, M_k^c \circ \tau \simeq M_k^b, M_k^b \circ \tau \simeq M_k^a, \\
W_k^a \circ \tau &\simeq W_k^c, W_k^c \circ \tau \simeq W_k^b, \text{ and } W_k^b \circ \tau \simeq W_k^a.
\end{aligned}$$

So $\{M_k^0\}$, $\{W_k^0\}$, $\{M_k^a, M_k^b, M_k^c\}$, and $\{W_k^a, W_k^b, W_k^c\}$ are G -stable sets. The automorphism τ of $V_{\sqrt{2}A_2}$ fixes ω^3 and so W_k^0 is invariant under τ . Hence we can take τ as $\phi(\tau, W_k^0)$ in Section 2.1. For these G -stable sets, we can take the 2-cocycles α in Section 2.1 to be trivial. Let $\xi = e^{2\pi\sqrt{-1}/3}$. We set $M_k^{0(i)} = \{v \in M_k^0 \mid \tau v = \xi^i v\}$ and $W_k^{0(i)} = \{v \in W_k^0 \mid \tau v = \xi^i v\}$ for $i \in \mathbb{Z}$. Note that $\mathcal{W} = M_k^{0(0)}$. By Theorem 1, we have $M_k^{0(i)}, W_k^{0(i)}$, ($i = 0, 1, 2$) M_k^a , and W_k^a are inequivalent irreducible $\mathcal{W}$ -modules. Moreover, $M_k^a \simeq M_k^b \simeq M_k^c$ and $W_k^a \simeq W_k^b \simeq W_k^c$ as $\mathcal{W}$ -modules. The contragredient modules of these $\mathcal{W}$ -modules are

$$\begin{aligned}
(M_k^{0(i)})' &\simeq M_k^{0(-i)}, (W_k^{0(i)})' \simeq W_k^{0(-i)}, \quad (i = 0, 1, 2), \\
(M_k^a)' &\simeq M_k^a \text{ and } (W_k^a)' \simeq W_k^a.
\end{aligned}$$

All the other irreducible $\mathcal{W}$ -modules occur as submodules of irreducible τ^i -twisted M_k^0 -modules for $i = 1, 2$.

4.3 An upper bound for the fusion rule

We review some notations and formulas for the Zhu algebra $A(V)$ of an arbitrary vertex operator algebra V in [14] and the $A(V)$ -bimodule $A(L)$

of an arbitrary V -module L in [10] and [13]. For $u, v \in V$ with u being homogeneous, define two bilinear operations

$$\begin{aligned} u * v &= \text{Res}_z(Y(u, z)v \frac{(1+z)^{\text{wt}u}}{z}), \\ u \circ v &= \text{Res}_z(Y(u, z)v \frac{(1+z)^{\text{wt}u}}{z^2}). \end{aligned}$$

We extend $*$ and $\circ$ for arbitrary $u, v \in V$ by linearity. Let $O(V)$ be the subspace of V spanned by all $u \circ v$ for $u \in V, v \in L$. Set $A(V) = V/O(V)$. By [14, Theorem 2.1.1.], $O(V)$ is a two-sided ideal with respect to the operation $*$ and $(A(V), *)$ is an associative algebra with identity $1 + O(V)$. For every V -module N , $N(0)$ is a left $A(V)$ -module.

Let L be a V -module. For $u \in V, v \in L$ with u being homogeneous, define three bilinear operations

$$\begin{aligned} u * v &= \text{Res}_z(Y(u, z)v \frac{(1+z)^{\text{wt}u}}{z}), \\ v * u &= \text{Res}_z(Y(u, z)v \frac{(1+z)^{\text{wt}u-1}}{z}), \\ u \circ v &= \text{Res}_z(Y(u, z)v \frac{(1+z)^{\text{wt}u}}{z^2}). \end{aligned}$$

We extend $*$ and $\circ$ for arbitrary $u \in V, v \in L$ by linearity. Let $O(L)$ be the subspace of L spanned by all $u \circ v$ for $u \in V, v \in L$. By [10, Theorem 1.5.1.], $O(L)$ is a two-sided ideal with respect to the operation $*$. Thus it induces an operation on $A(L) = L/O(L)$. Denote by $[v]$ the image of $v \in L$ in $A(L)$. $A(L)$ is a $A(V)$ -bimodule under the operation $*$. Using $A(L)$, we have an upper bound for every fusion rule.

Lemma 5. ([13] Proposition 2.10.) *Let $L^i = \bigoplus_{n=0}^{\infty} L^i(n)$ ($i = 1, 2, 3$) be irreducible V -modules. Then*

$$\dim_{\mathbb{C}} I_V \begin{pmatrix} L^3 \\ L^1 \quad L^2 \end{pmatrix} \leq \dim_{\mathbb{C}} \text{Hom}_{\mathbb{C}} \left(L^3(0)^* \otimes_{A(V)} A(L^1) \otimes_{A(V)} L^2(0) \right).$$

4.4 The fusion rules for irreducible $\mathcal{W}$ -modules which occur as submodules of irreducible M_k^0 -modules

Let L^i ($i = 1, 2, 3$) be irreducible $\mathcal{W}$ -modules such that L^1 and L^2 occur as submodules of irreducible M_k^0 -modules. In this subsection we determine the

fusion rule of type $\binom{L^3}{L^1 L^2}$. As a first step, we give a lower bound for every fusion rule by using Theorem 2.

Lemma 6. (1) *The fusion rule of following types is greater than or equal to 1: Let $i, j \in \{0, 1, 2\}$.*

$$\begin{aligned} & \binom{M_k^{0(i+j)}}{M_k^{0(i)} M_k^{0(j)}}, \quad \binom{W_k^{0(i+j)}}{W_k^{0(i)} W_k^{0(j)}}, \quad \binom{W_k^{0(i+j)}}{M_k^{0(i)} W_k^{0(j)}}, \quad \binom{M_k^a}{M_k^{0(i)} M_k^a}, \\ & \binom{W_k^a}{M_k^{0(i)} W_k^a}, \quad \binom{M_k^a}{W_k^{0(i)} W_k^a}, \quad \binom{W_k^a}{W_k^{0(i)} W_k^a}. \end{aligned}$$

(2) *The fusion rule of following types is greater than or equal to 2:*

$$\binom{M_k^a}{M_k^a M_k^a}, \quad \binom{W_k^a}{M_k^a W_k^a}, \quad \binom{W_k^a}{W_k^a W_k^a}.$$

Proof. We use notations in Section 3. The fusion rules for M_k^0 are obtained in [12].

- (1) We consider the case that $\mathcal{S}_1 = \mathcal{S}_2 = \mathcal{S}_3 = \{W_k^0\}$. We have $\dim_{\mathbb{C}} I_{M_k^0} \left(\binom{W_k^0}{W_k^0 W_k^0} \right) = 1$. Let $\mathbb{C}x_i$ ($i = 0, 1, 2$) be the one dimensional G -modules such that $\tau \cdot x_i = \xi^i x_i$. Fix nonzero $v^i \in \text{Hom}_{\mathbb{C}G}(\mathbb{C}x_i, W_k^{0(i)})$. Then

$$\begin{aligned} W_k^0 &= \bigoplus_{i=0}^2 x_i \otimes \text{Hom}_{\mathbb{C}G}(\mathbb{C}x_i, W_k^{0(i)}), \\ \mathcal{I} &= I_{M_k^0} \left(\binom{W_k^0}{W_k^0 W_k^0} \right) \otimes_{\mathbb{C}} W_k^0 \otimes_{\mathbb{C}} W_k^0, \text{ and} \\ \mathcal{I}_{W_k^{0(i)}, W_k^{0(j)}}(v^i, v^j) &= I_{M_k^0} \left(\binom{W_k^0}{W_k^0 W_k^0} \right) \otimes (x_i \otimes v^i) \otimes (x_j \otimes v^j). \end{aligned}$$

Let $f \in I_{M_k^0} \left(\binom{W_k^0}{W_k^0 W_k^0} \right)$ be a nonzero intertwining operator. Then $\tau f = f$ by the construction of intertwining operators in [12]. So

$$(\tau \otimes e(W_k^0)) \cdot f \otimes (x_i \otimes v^i) \otimes (x_j \otimes v^j) = \xi^{i+j} f \otimes (x_i \otimes v^i) \otimes (x_j \otimes v^j)$$

and $\dim_{\mathbb{C}} I_{\mathcal{W}} \left(\binom{W_k^{0(i+j)}}{W_k^{0(i)} W_k^{0(j)}} \right) \geq 1$. We can compute the other cases in the same way.

- (2) We consider the case that $\mathcal{S}_1 = \mathcal{S}_2 = \mathcal{S}_3 = \{M_k^a, M_k^b, M_k^c\}$. For $i_1, i_2, i_3 \in \{a, b, c\}$, we have

$$\dim_{\mathbb{C}} I_{M_k^0} \begin{pmatrix} M_k^{i_3} \\ M_k^{i_1} & M_k^{i_2} \end{pmatrix} = \begin{cases} 1, & \text{if } \{i_1, i_2, i_3\} = \{a, b, c\}, \\ 0, & \text{otherwise.} \end{cases}$$

We define an action of τ on $\{a, b, c\}$ by $\tau(a) = b, \tau(b) = c$ and $\tau(c) = a$. It is possible to take $\phi(\tau, M_k^i)$ ($i = a, b, c$) such that

$$\phi(\tau, M_k^c) \phi(\tau, M_k^b) \phi(\tau, M_k^a) = \text{id}_{M_k^a}.$$

Fix a nonzero $v^a \in M_k^a$ and set $v^b = \phi(\tau, M_k^a) v^a \in M_k^b, v^c = \phi(\tau, M_k^b) v^b \in M_k^c$. Fix nonzero $f_{a,b,c} \in I_{M_k^0} \begin{pmatrix} M_k^c \\ M_k^a & M_k^b \end{pmatrix}, f_{b,a,c} \in I_{M_k^0} \begin{pmatrix} M_k^c \\ M_k^b & M_k^a \end{pmatrix}$ and set

$$\begin{aligned} f_{b,c,a} &= \tau f_{a,b,c} \in I_{M_k^0} \begin{pmatrix} M_k^a \\ M_k^b & M_k^c \end{pmatrix}, \\ f_{c,a,b} &= \tau f_{b,c,a} \in I_{M_k^0} \begin{pmatrix} M_k^b \\ M_k^c & M_k^a \end{pmatrix}, \\ f_{c,b,a} &= \tau f_{b,a,c} \in I_{M_k^0} \begin{pmatrix} M_k^a \\ M_k^c & M_k^b \end{pmatrix}, \\ f_{a,c,b} &= \tau f_{c,b,a} \in I_{M_k^0} \begin{pmatrix} M_k^b \\ M_k^a & M_k^c \end{pmatrix}. \end{aligned}$$

We have

$$\mathcal{I} = \bigoplus_{\substack{i_1, i_2, i_3 \in \{a, b, c\} \\ \{i_1, i_2, i_3\} = \{a, b, c\}}} I_{M_k^0} \begin{pmatrix} M_k^{i_3} \\ M_k^{i_1} & M_k^{i_2} \end{pmatrix} \otimes_{\mathbb{C}} M_k^{i_1} \otimes_{\mathbb{C}} M_k^{i_2}$$

and

$$\begin{aligned} \mathcal{I}_{M_k^a, M_k^a} &= \bigoplus_{i=0,1,2} \mathbb{C} f_{\tau^i(a), \tau^i(b), \tau^i(c)} \otimes v^{\tau^i(a)} \otimes v^{\tau^i(b)} \\ &\quad \oplus \bigoplus_{i=0,1,2} \mathbb{C} f_{\tau^i(b), \tau^i(a), \tau^i(c)} \otimes v^{\tau^i(b)} \otimes v^{\tau^i(a)}. \end{aligned}$$

Since $\bigoplus_{i=0,1,2} \mathbb{C} f_{\tau^i(a), \tau^i(b), \tau^i(c)} \otimes v^{\tau^i(a)} \otimes v^{\tau^i(b)}$ and $\bigoplus_{i=0,1,2} \mathbb{C} f_{\tau^i(b), \tau^i(a), \tau^i(c)} \otimes v^{\tau^i(b)} \otimes v^{\tau^i(a)}$ are isomorphic irreducible $\mathcal{A}_{\alpha_3}(G, \mathcal{S}_3)$ -modules, we have $\dim_{\mathbb{C}} I_{\mathcal{W}} \begin{pmatrix} M_k^a \\ M_k^a & M_k^a \end{pmatrix} \geq 2$. We can compute the other cases in the same way.

□

We will show that every fusion rule meets the lower bound obtained in Lemma 6. We use Lemma 5. For an irreducible $\mathcal{W}$ -module $N = \bigoplus_{n=0}^{\infty} N(n)$, we fix a nonzero vector in $N(0)$ and denote it by w_N . By the same argument as in [3, Lemma 5.2], N is spanned by the vectors of the form

$$L(-m_1) \cdots L(-m_p) J(-n_1) \cdots J(-n_q) w_N \quad (5)$$

with $m_1 \geq \cdots \geq m_p \geq 1, n_1 \geq \cdots \geq n_q \geq 1, p = 0, 1, \dots$, and $q = 0, 1, \dots$. By the same argument as in [3, Section 5.4], we have for $n \leq -1$ and $u \in N$,

$$\begin{aligned} L(n)u &= (-1)^{n-1}[\omega] * [u] + (-1)^{n-1}n[u] * [\omega] + (-1)^n \text{wt} u [u], \\ J(n)u &= (-1)^n \left(n[J(-1)u] + (n+1)[J[0]u] - (n+1)[J] * [u] \right. \\ &\quad \left. - \frac{n(n+1)}{2}[u] * [J] \right) \end{aligned} \quad (6)$$

in $A(N)$. Using formula (6) and commutation relations (4) repeatedly, it is shown that $A(N)$ is generated by $\{J(-1)^i w_N\}_{i=0}^{\infty}$ as a $A(\mathcal{W})$ -bimodule.

A *singular vector* w of weight h for N is by definition a vector w which satisfies $L(0)w = hw$ and $L(n)w = J(n)w = 0$ for $n \geq 1$. By commutation relations (4), it is easy to show that w is a singular vector of weight h if and only if $L(0)w = hw$ and $L(1)w = L(2)w = J(1)w = 0$. If $h - \text{wt} N(0) > 0$, then the submodule of N generated by w does not contain $N(0)$. We hence have $w = 0$.

Theorem 4. *The fusion rule $L^1 \times L^2$ is given by the following list, where $\{L^1, L^2\}$ is an arbitrary pair of irreducible $\mathcal{W}$ -modules which occur as sub-*

modules of irreducible M_k^0 -modules: Let $i, j \in \{0, 1, 2\}$.

$$\begin{aligned}
M_k^{0(i)} \times M_k^{0(j)} &= M_k^{0(i+j)}, \\
M_k^{0(i)} \times W_k^{0(j)} &= W_k^{0(i+j)}, \\
W_k^{0(i)} \times W_k^{0(j)} &= M_k^{0(i+j)} + W_k^{0(i+j)}, \\
M_k^{0(i)} \times M_k^a &= M_k^a, \\
M_k^{0(i)} \times W_k^a &= W_k^a, \\
W_k^{0(i)} \times M_k^a &= W_k^a, \\
W_k^{0(i)} \times W_k^a &= M_k^a + W_k^a, \\
M_k^a \times M_k^a &= \sum_{i=0}^2 M_k^{0(i)} + 2M_k^a, \\
M_k^a \times W_k^a &= \sum_{i=0}^2 W_k^{0(i)} + 2W_k^a, \\
W_k^a \times W_k^a &= \sum_{i=0}^2 M_k^{0(i)} + \sum_{i=0}^2 W_k^{0(i)} + 2M_k^a + 2W_k^a.
\end{aligned}$$

Proof. For an irreducible $\mathcal{W}$ -module N , h_N denotes the eigenvalue for $L(0)$ on $N(0)$ and k_N denotes the eigenvalue for $J(0)$ on $N(0)$. For all irreducible $\mathcal{W}$ -modules those eigenvalues are computed in [3]. For irreducible $\mathcal{W}$ -modules L^i ($i = 1, 2, 3$), set

$$F(L^1, L^2, L^3) = \dim_{\mathbb{C}} (L^3(0)^* \otimes_{A(\mathcal{W})} A(L^1) \otimes_{A(\mathcal{W})} L^2(0)).$$

Note that $F(L^1, L^2, L^3)$ is lower than or equal to the number of generators of $A(L^1)$ as a $A(\mathcal{W})$ -bimodule since the dimension of top level of every irreducible $\mathcal{W}$ -module is 1.

The simplicity of $\mathcal{W}$ implies that

$$\dim_{\mathbb{C}} I_{\mathcal{W}} \begin{pmatrix} L^3 \\ \mathcal{W} \ L^2 \end{pmatrix} = \begin{cases} 1, & \text{if } L^2 \simeq L^3, \\ 0, & \text{otherwise} \end{cases}$$

for all irreducible $\mathcal{W}$ -modules L^2 and L^3 . We consider the case $L^1 \neq \mathcal{W}$. We will show that $F(L^1, L^2, L^3)$ is lower than or equal to the lower bound for $\dim_{\mathbb{C}} I_{\mathcal{W}} \begin{pmatrix} L^3 \\ L^1 \ L^2 \end{pmatrix}$ given in Lemma 6. Then, by Lemma 3 and Lemma 5 we get the desired results. We use a computer algebra system Risa/Asir to find

singular vectors used in the following argument. The explicit forms of those singular vectors are given in Appendix.

- (1) We consider the case $L^1 = W_k^{0(1)}$. Since $(5\sqrt{-3}L(-1) + J(-1))w_{W_k^{0(1)}}$ is a singular vector and $A(W_k^{0(1)})$ is generated by $\{J(-1)^i w_{W_k^{0(1)}}\}_{i=0}^\infty$ as a $A(\mathcal{W})$ -bimodule, $A(W_k^{0(1)})$ is generated by $w_{W_k^{0(1)}}$ as a $A(\mathcal{W})$ -bimodule. So $F(W_k^{0(1)}, L^2, L^3) \leq 1$. Set

$$\begin{aligned} w_1 = & \left(-30\sqrt{-3}L(-1)J(-1) + 39\sqrt{-3}J(-2) \right. \\ & \left. + 5J(-1)^2 + 336L(-2) + 405L(-1)^2 \right) w_{W_k^{0(1)}}. \end{aligned}$$

Since w_1 is a singular vector, we have a relation

$$\begin{aligned} 0 = & 50\sqrt{-3}([\omega]^2 * [w_{W_k^{0(1)}}] + [w_{W_k^{0(1)}}] * [\omega]^2) \\ & - 20\sqrt{-3}([\omega] * [w_{W_k^{0(1)}}] + [w_{W_k^{0(1)}}] * [\omega]) \\ & + 4\sqrt{-3}[w_{W_k^{0(1)}}] - 100\sqrt{-3}[\omega] * [w_{W_k^{0(1)}}] * [\omega] \\ & - 5[w_{W_k^{0(1)}}] * [J] + 5[J] * [w_{W_k^{0(1)}}] \end{aligned}$$

in $A(W_0^k(1))$ by using formulas (6). Therefore,

$$\begin{aligned} 0 = & (50\sqrt{-3}(h_{L^2}^2 + h_{L^3}^2) - 20\sqrt{-3}(h_{L^2} + h_{L^3}) \\ & + 4\sqrt{-3} - 100\sqrt{-3}h_{L^2}h_{L^3} - 5k_{L^2} + 5k_{L^3}) w_{(L^3)'} \otimes [w_{W_k^{0(1)}}] \otimes w_{L^2} \end{aligned}$$

in $L^3(0)^* \otimes_{A(\mathcal{W})} A(W_k^{0(1)}) \otimes_{A(\mathcal{W})} L^2(0)$. Set

$$\begin{aligned} \psi(h_{L^2}, k_{L^2}, h_{L^2}, k_{L^3}) = & 50\sqrt{-3}(h_{L^2}^2 + h_{L^3}^2) - 20\sqrt{-3}(h_{L^2} + h_{L^3}) \\ & + 4\sqrt{-3} - 100\sqrt{-3}h_{L^2}h_{L^3} - 5k_{L^2} + 5k_{L^3}. \end{aligned}$$

If $\psi(h_{L^2}, k_{L^2}, h_{L^2}, k_{L^3}) \neq 0$, then $w_{(L^3)'} \otimes [w_{W_k^{0(1)}}] \otimes w_{L^2} = 0$ and $F(W_k^{0(1)}, L^2, L^3) = 0$. By computing $\psi(h_{L^2}, k_{L^2}, h_{L^3}, k_{L^3})$ for all pairs (L^2, L^3) of 20 irreducible $\mathcal{W}$ -modules, we have $F(W_k^{0(1)}, L^2, L^3) \leq 1$ if the pair (L^2, L^3) is one of

$$\begin{aligned} & (M_k^{0(i)}, W_k^{0(1+i)}), (W_k^{0(i)}, M_k^{0(1+i)}), (W_k^{0(i)}, W_k^{0(1+i)}), \quad (i = 0, 1, 2), \\ & (M_k^a, W_k^a), (W_k^a, M_k^a), \text{ and } (W_k^a, W_k^a) \end{aligned}$$

and $F(W_k^{0(1)}, L^2, L^3) = 0$ otherwise. Combining these results and Lemma 6, we have

$$\begin{aligned} W_k^{0(1)} \times M_k^{0(i)} &= W_k^{0(1+i)}, \\ W_k^{0(1)} \times W_k^{0(i)} &= M_k^{0(1+i)} + W_k^{0(1+i)}, \\ W_k^{0(1)} \times M_k^a &= W_k^a, \\ W_k^{0(1)} \times W_k^a &= M_k^a + W_k^a. \end{aligned}$$

In the case $L^1 = W_k^{0(2)}$, we can compute the fusion rules in the same way. In the case $L^1 = M_k^{0(i)}$ ($i = 1, 2$), it is shown that $A(M_k^{0(i)})$ is generated by $[w_{M_k^{0(i)}}]$ in the same way. But in these cases we need two singular vectors $v_{31} \in M_k^{0(i)}(3)$ and $v_{41} \in M_k^{0(i)}(4)$. We can compute the fusion rules by using two relations $v_{41} = 0$ and $J(-1)v_{31} = 0$.

- (2) We consider the case $L^1 = W_k^{0(0)}$. There are two singular vectors $u_{2i} \in W_k^{0(0)}(2)$ ($i = 1, 2$) and there is one singular vector $u_{41} \in W_k^{0(0)}(4)$. Since $u_{21} = 0$, $A(W_k^{0(0)})$ is generated by $\{w_{W_k^{0(0)}}, J(-1)w_{W_k^{0(0)}}\}$ as a $A(\mathcal{W})$ -bimodule and for $X \in W_k^{0(0)}$ we have $a_1(L^2, L^3; X), a_2(L^2, L^3; X) \in \mathbb{C}$ such that

$$\begin{aligned} w_{(L^3)'} \otimes [X] \otimes w_{L^2} &= a_1(L^2, L^3; X)w_{(L^3)'} \otimes [w_{W_k^{0(0)}}] \otimes w_{L^2} \\ &\quad + a_2(L^2, L^3; X)w_{(L^3)'} \otimes [J(-1)w_{W_k^{0(0)}}] \otimes w_{L^2}. \end{aligned}$$

Therefore, $F(W_k^{0(0)}, L^2, L^3) \leq 2$. Set a matrix

$$A = \begin{pmatrix} a_1(L^2, L^3; u_{22}) & a_2(L^2, L^3; u_{22}) \\ a_1(L^2, L^3; J(-1)u_{22}) & a_2(L^2, L^3; J(-1)u_{22}) \\ a_1(L^2, L^3; v_{41}) & a_2(L^2, L^3; v_{41}) \\ a_1(L^2, L^3; J(-1)u_{41}) & a_2(L^2, L^3; J(-1)u_{41}) \end{pmatrix}.$$

We have

$$A \begin{pmatrix} (w_{(L^3)'} \otimes [w_{W_k^{0(0)}}] \otimes w_{L^2}) \\ (w_{(L^3)'} \otimes [J(-1)w_{W_k^{0(0)}}] \otimes w_{L^2}) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}.$$

If $\text{rank} A = 2$, then $w_{(L^3)'} \otimes [w_{W_k^{0(0)}}] \otimes w_{L^2} = w_{(L^3)'} \otimes [J(-1)w_{W_k^{0(0)}}] \otimes w_{L^2} = 0$ in $L^3(0)^* \otimes_{A(\mathcal{W})} A(W_k^{0(0)}) \otimes_{A(\mathcal{W})} L^2(0)$. If $\text{rank} A = 1$, then $w_{(L^3)'} \otimes [w_{W_k^{0(0)}}] \otimes w_{L^2}$ and $w_{(L^3)'} \otimes [J(-1)w_{W_k^{0(0)}}] \otimes w_{L^2}$ are linearly dependent. We hence have

$$F(W_k^{0(0)}, L^2, L^3) \leq \begin{cases} 0, & \text{if } \text{rank} A = 2, \\ 1, & \text{if } \text{rank} A = 1, \\ 2, & \text{if } \text{rank} A = 0. \end{cases}$$

By computing the rank of A for all pairs (L^2, L^3) of 20 irreducible $\mathcal{W}$ -modules, we have for $i = 0, 1, 2$

$$\begin{aligned} W_k^{0(0)} \times M_k^{0(i)} &= W_k^{0(i)}, \\ W_k^{0(0)} \times W_k^{0(i)} &= M_k^{0(i)} + W_k^{0(i)}, \\ W_k^{0(0)} \times M_k^a &= W_k^a, \\ W_k^{0(0)} \times W_k^a &= M_k^a + W_k^a. \end{aligned}$$

In the case that L^1 is one of M_k^a and W_k^a , we can compute the fusion rules in the same way. We roughly explain each case. In the case $L^1 = M_k^a$, there are two singular vectors v_{21}, v_{22} in $M_k^a(2)$ and three singular vectors v_{61}, v_{62}, v_{63} in $M_k^a(6)$. Set the same matrix for $v_{22}, v_{61}, v_{62}, v_{63}, J(-1)v_{61}, J(-1)v_{62}$ and $J(-1)v_{63}$ as in the case of $W_k^{0(0)}$. By computing the rank of the matrix, we can determine the fusion rules. In the case $L^1 = W_k^a$, there is a singular vector v_{21} in $W_k^a(2)$ and there are two singular vectors v_{41}, v_{42} in $W_k^a(4)$. Set the same matrix for $v_{41}, v_{42}, J(-1)v_{41}$ and $J(-1)v_{42}$ as in the case of $W_k^{0(0)}$. By computing the rank of the matrix, we can determine the fusion rules.

□

5 Appendix

We give some singular vectors in irreducible $\mathcal{W}$ -modules used in Theorem 4. We omit $w_N \in N(0)$ from the explicit form of every singular vector in a $\mathcal{W}$ -module N .

• $M_k^{0(1)}$.

- (1) $9\sqrt{-3}L(-1) + J(-1),$
- (2) $6255L(-1)J(-2) - 375\sqrt{-3}L(-1)J(-1)^2 + 36960L(-2)J(-1)$
 $+ 5175L(-1)^2J(-1) - 26208J(-3) + 1425\sqrt{-3}J(-2)J(-1)$
 $+ 25J(-1)^3 + 23040\sqrt{-3}L(-3) + 147600\sqrt{-3}L(-2)L(-1)$
 $- 44625\sqrt{-3}L(-1)^3,$
- (3) $-36720\sqrt{-3}L(-1)J(-3) + 324L(-1)J(-2)J(-1)$
 $- 16\sqrt{-3}L(-1)J(-1)^3 + 9360\sqrt{-3}L(-2)J(-2)$
 $+ 12852\sqrt{-3}L(-1)^2J(-2) + 2400L(-2)J(-1)^2$
 $+ 462L(-1)^2J(-1)^2 - 5040\sqrt{-3}L(-3)J(-1)$
 $+ 8640\sqrt{-3}L(-2)L(-1)J(-1) - 5232\sqrt{-3}L(-1)^3J(-1)$
 $+ 35280\sqrt{-3}J(-4) - 819J(-2)^2$
 $+ 76\sqrt{-3}J(-2)J(-1)^2 + J(-1)^4$
 $- 751680L(-4) - 1028160L(-3)L(-1)$
 $+ 254880L(-2)L(-1)^2 + 16929L(-1)^4.$

We obtain singular vectors in $M_k^{0(2)}$ by replacing $J(n)$ with $-J(n)$ in the above vectors.

• $W_k^{0(0)}$.

- (1) $-70\sqrt{-3}L(-1)J(-1) + 91\sqrt{-3}J(-2) - 5J(-1)^2 - 2496L(-2) + 195L(-1)^2,$
- (2) $-70\sqrt{-3}L(-1)J(-1) + 91\sqrt{-3}J(-2) + 5J(-1)^2 + 2496L(-2) - 195L(-1)^2,$
- (3) $-1500L(-1)J(-2)J(-1) + 1200L(-2)J(-1)^2 + 750L(-1)^2J(-1)^2$
 $+ 3600J(-3)J(-1) + 825J(-2)^2 + J(-1)^4$
 $- 633600L(-4) + 46800L(-3)L(-1) + 230400L(-2)^2$
 $- 126000L(-2)L(-1)^2 + 50625L(-1)^4,$

• $W_k^{0(1)}$.

- (1) $5\sqrt{-3}L(-1) + J(-1),$
- (2) $-30\sqrt{-3}L(-1)J(-1) + 39\sqrt{-3}J(-2) + 5J(-1)^2 + 336L(-2) + 405L(-1)^2.$

We obtain singular vectors in $W_k^{0(2)}$ by replacing $J(n)$ with $-J(n)$ in the above vectors.

• W_k^a .

- (1) $J(-1)^2 - 30L(-2) + 75L(-1)^2$,
- (2) $-5040\sqrt{-3}L(-1)J(-3) - 4980L(-1)J(-2)J(-1)$
 $+ 200\sqrt{-3}L(-1)J(-1)^3 + 7668\sqrt{-3}L(-2)J(-2)$
 $+ 990\sqrt{-3}L(-1)^2J(-2) + 9660L(-2)J(-1)^2$
 $+ 7350L(-1)^2J(-1)^2 + 35760\sqrt{-3}L(-3)J(-1)$
 $- 78960\sqrt{-3}L(-2)L(-1)J(-1) + 34200\sqrt{-3}L(-1)^3J(-1)$
 $+ 5208\sqrt{-3}J(-4) + 12780J(-3)J(-1)$
 $+ 2622J(-2)J(-2) - 310\sqrt{-3}J(-2)J(-1)^2$
 $+ 25J(-1)^4 - 9000L(-4)$
 $- 255780L(-3)L(-1) - 28044L(-2)^2$
 $+ 557460L(-2)L(-1)^2 - 78975L(-1)^4$,
- (3) $5040\sqrt{-3}L(-1)J(-3) - 4980L(-1)J(-2)J(-1)$
 $- 200\sqrt{-3}L(-1)J(-1)^3 - 7668\sqrt{-3}L(-2)J(-2)$
 $- 990\sqrt{-3}L(-1)^2J(-2) + 9660L(-2)J(-1)^2$
 $+ 7350L(-1)^2J(-1)^2 - 35760\sqrt{-3}L(-3)J(-1)$
 $+ 78960\sqrt{-3}L(-2)L(-1)J(-1) - 34200\sqrt{-3}L(-1)^3J(-1)$
 $- 5208\sqrt{-3}J(-4) + 12780J(-3)J(-1)$
 $+ 2622J(-2)J(-2) + 310\sqrt{-3}J(-2)J(-1)^2$
 $+ 25J(-1)^4 - 9000L(-4)$
 $- 255780L(-3)L(-1) - 28044L(-2)^2$
 $+ 557460L(-2)L(-1)^2 - 78975L(-1)^4$.

• M_k^a .

- (1) $8\sqrt{-3}L(-1)J(-1) - 6\sqrt{-3}J(-2) + J(-1)^2 + 90L(-2) + 27L(-1)^2$,
- (2) $-8\sqrt{-3}L(-1)J(-1) + 6\sqrt{-3}J(-2) + J(-1)^2 + 90L(-2) + 27L(-1)^2$,

There are another three singular vectors v_{6i} ($i = 1, 2, 3$) in M_k^a ,

$$\begin{aligned}
v_{61} = & 63987840\sqrt{-3}L(-1)J(-5) - 1059480L(-1)J(-4)J(-1) \\
& + 1587600L(-1)J(-3)J(-2) + 14400\sqrt{-3}L(-1)J(-3)J(-1)^2 \\
& - 26208\sqrt{-3}L(-1)J(-2)^2J(-1) + 4260L(-1)J(-2)J(-1)^3 \\
& + 32\sqrt{-3}L(-1)J(-1)^5 + 23284800\sqrt{-3}L(-2)J(-4) \\
& - 19867680\sqrt{-3}L(-1)^2J(-4) + 145800L(-2)J(-3)J(-1) \\
& - 1414260L(-1)^2J(-3)J(-1) - 427140L(-2)J(-2)^2 \\
& - 155358L(-1)^2J(-2)^2 - 15840\sqrt{-3}L(-2)J(-2)J(-1)^2 \\
& + 10224\sqrt{-3}L(-1)^2J(-2)J(-1)^2 + 3990L(-2)J(-1)^4 \\
& - 447L(-1)^2J(-1)^4 - 25401600\sqrt{-3}L(-3)J(-3) \\
& - 5443200\sqrt{-3}L(-2)L(-1)J(-3) + 1995840\sqrt{-3}L(-1)^3J(-3) \\
& - 659880L(-3)J(-2)J(-1) + 1666440L(-2)L(-1)J(-2)J(-1) \\
& + 341388L(-1)^3J(-2)J(-1) - 41280\sqrt{-3}L(-3)J(-1)^3 \\
& + 32640\sqrt{-3}L(-2)L(-1)J(-1)^3 + 7872\sqrt{-3}L(-1)^3J(-1)^3 \\
& + 4294080\sqrt{-3}L(-4)J(-2) + 16420320\sqrt{-3}L(-3)L(-1)J(-2) \\
& + 4989600\sqrt{-3}L(-2)^2J(-2) - 1360800\sqrt{-3}L(-2)L(-1)^2J(-2) \\
& + 122472\sqrt{-3}L(-1)^4J(-2) - 3483720L(-4)J(-1)^2 \\
& - 1742220L(-3)L(-1)J(-1)^2 + 38700L(-2)^2J(-1)^2 \\
& + 588420L(-2)L(-1)^2J(-1)^2 - 169749L(-1)^4J(-1)^2 \\
& - 42370560\sqrt{-3}L(-5)J(-1) - 45861120\sqrt{-3}L(-4)L(-1)J(-1) \\
& - 2332800\sqrt{-3}L(-3)L(-2)J(-1) - 31164480\sqrt{-3}L(-3)L(-1)^2J(-1) \\
& - 3542400\sqrt{-3}L(-2)^2L(-1)J(-1) + 4429440\sqrt{-3}L(-2)L(-1)^3J(-1) \\
& + 23328\sqrt{-3}L(-1)^5J(-1) - 101787840\sqrt{-3}J(-6) \\
& - 546480J(-5)J(-1) - 1186920J(-4)J(-2) \\
& + 5280\sqrt{-3}J(-4)J(-1)^2 - 2381400J(-3)^2 \\
& - 47520\sqrt{-3}J(-3)J(-2)J(-1) - 3420J(-3)J(-1)^3 \\
& + 11088\sqrt{-3}J(-2)^3 + 870J(-2)^2J(-1)^2 \\
& - 88\sqrt{-3}J(-2)J(-1)^4 + J(-1)^6 \\
& + 879076800L(-6) + 990072720L(-5)L(-1) \\
& + 65091600L(-4)L(-2) + 323666280L(-4)L(-1)^2 \\
& + 102173400L(-3)^2 + 73823400L(-3)L(-2)L(-1) \\
& - 3027780L(-3)L(-1)^3 - 152523000L(-2)^3 \\
& + 95652900L(-2)^2L(-1)^2 - 15326010L(-2)L(-1)^4 \\
& - 505197L(-1)^6.
\end{aligned}$$

The vector v_{62} is obtained by replacing $J(n)$ with $-J(n)$ in v_{61} .

$$\begin{aligned}
v_{63} = & -1478136600L(-1)J(-4)J(-1) + 423979920L(-1)J(-3)J(-2) \\
& - 1273500L(-1)J(-2)J(-1)^3 - 29005560L(-2)J(-3)J(-1) \\
& + 58538700L(-1)^2J(-3)J(-1) + 134322300L(-2)J(-2)^2 \\
& - 133840350L(-1)^2J(-2)^2 + 1341750L(-2)J(-1)^4 \\
& + 680625L(-1)^2J(-1)^4 - 778588200L(-3)J(-2)J(-1) \\
& + 143310600L(-2)L(-1)J(-2)J(-1) + 37975500L(-1)^3J(-2)J(-1) \\
& - 2232505800L(-4)J(-1)^2 - 311863500L(-3)L(-1)J(-1)^2 \\
& + 586133100L(-2)^2J(-1)^2 - 239341500L(-2)L(-1)^2J(-1)^2 \\
& + 127186875L(-1)^4J(-1)^2 - 393666480J(-5)J(-1) \\
& - 1236672360J(-4)J(-2) - 104786136J(-3)^2 \\
& + 3968100J(-3)J(-1)^3 + 2139750J(-2)^2J(-1)^2 \\
& + 625J(-1)^6 + 275225065920L(-6) \\
& + 450006898320L(-5)L(-1) + 221829042960L(-4)L(-2) \\
& - 60223224600L(-4)L(-1)^2 + 121440173400L(-3)^2 \\
& + 145205865000L(-3)L(-2)L(-1) - 22975690500L(-3)L(-1)^3 \\
& - 5826151800L(-2)^3 + 65516566500L(-2)^2L(-1)^2 \\
& - 55664516250L(-2)L(-1)^4 + 5974171875L(-1)^6.
\end{aligned}$$

References

- [1] R. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, *Proc. Nat. Acad. Sci. U.S.A.* **83** (1986), 3068–3071.
- [2] R. Dijkgraaf, C. Vafa, E. Verlinde, and H. Verlinde, The operator algebra of orbifold models, *Comm. Math. Phys.* **123** (1989), 485–526.
- [3] C. Dong, C.H. Lam, K. Tanabe, H. Yamada, and K. Yokoyama, $\mathbb{Z}_3$ symmetry and W_3 algebra in lattice vertex operator algebras, *Pacific J. Math.* **215** (2004), 245–296.
- [4] C. Dong and J. Lepowsky, Generalized Vertex Algebras and Relative Vertex Operators, Progress in Math., Vol. **112**, Birkhauser, Boston, 1993.
- [5] C. Dong and G. Mason, On quantum Galois theory, *Duke Math. J.* **86** (1997), 305–321.

- [6] C. Dong and G. Yamskulna, Vertex operator algebras, generalized doubles and dual pairs, *Math. Z.* **241** (2002), 397–423.
- [7] V. A. Fateev and A. B. Zamolodchikov, Conformal quantum field theory models in two dimensions having $\mathbb{Z}_3$ symmetry, *Nuclear Physics* **B280** (1987), 644–660.
- [8] I. Frenkel, Y. Huang, and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, *Mem. Amer. Math. Soc.* **104** (1993).
- [9] I. B. Frenkel, J. Lepowsky and A. Meurman, *Vertex Operator Algebras and the Monster*, Pure and Applied Math., Vol. **134**, Academic Press, 1988.
- [10] I. Frenkel and Y. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, *Duke Math. J.* **66** (1992), 123–168.
- [11] K. Kitazume, C. Lam and H. Yamada, 3-state Potts model, moonshine vertex operator algebra and $3A$ elements of the monster group, *IMRS*, **23** (2003), 1269–1303.
- [12] C. Lam and H. Yamada, $\mathbb{Z}_2 \times \mathbb{Z}_2$ codes and vertex operator algebras, *J. Algebra* **224** (2000), 268–291.
- [13] H. Li, Determining fusion rules by $A(V)$ -modules and bimodules, *J. Algebra* **212** (1999), 515–556.
- [14] Y. Zhu, Modular invariance of characters of vertex operator algebras, *J. Amer. Math. Soc.* **9** (1996), 237–302.